



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Mr. Palmer is right, sometimes he is wrong, and sometimes he is just 'betwixt and between.' Take the case of the genus whose name, in two forms, heads this paragraph. Mr. Palmer says truly that Vieillot wrote the word *Thriothorus*; for so it appears on p. 45 of my well-thumbed copy of his 'Analyse,' 1816, though this is the page which the A. O. U. Committee cite for *Thryothorus*. But if Mr. Palmer had looked further into Vieillot's 'Analyse,' he would have found *Thryothorus* tucked away near the bottom of the right hand column of p. 70, in an alphabetical list of the new genera of the book, where the etymologies are given. There the etymon of the first element of the word is stated as '*θρίον, juncus*'; and as the correct form resulting is *Thryothorus*, I think the Committee can defend their use of it, though they may have to cite p. 70 instead of p. 45 for it. At the same time, it offers a nice case for hair-splitting; for the previous *Thriothorus* of p. 45 cannot be brushed aside as a "typographical error," since Vieillot makes his intention clear by there writing 'THRIOTHORE, *Thriothorus*.' I commend the case to nomenclatural casuists.

While on the genus *Thryothorus*, I may inquire further how it happens that we have changed the name of *T. bewickii leucogaster* (Baird, 1864) to *T. b. bairdi* (Ridgway, 1885). The fact that there is a *Troglodytes leucogastra*, Gould, P. Z. S., 1836, p. 89 (which Baird mistook for the subspecies of *Thryothorus bewickii* which he named *leucogaster* in Rev. A. B., 1864, p. 127) does not affect the case one way or another. Gould's bird is now *Uropsila leucogastra*; it is also *Cyphorhinus pusillus* of Sclater, *Heterhina pusilla* of Baird, etc., and this cannot outlaw the use of the name *leucogaster* in the genus *Thryothorus*. Bewick's Wren has never been referred to the genus *Troglodytes* since one of its subspecies was called *leucogaster*, and of course there is no rule of nomenclature, express or implied, which requires us to change a specific name in one genus for the reason that the same has been used in another genus. Parity of reasoning—or rather, of unreasoning—would require us to reject *Cistothorus marianæ*, because there was a prior *Troglodytes marianæ*. I submit, therefore, that *Thryothorus bewickii bairdi*, No. 719b of the A. O. U. Lists, 1886 and 1895, should stand as *T. b. leucogaster*.—ELLIOT COUES, *Washington, D. C.*

The Hudsonian Chickadee breeding in Southern Vermont.—On June 29, 1895, I found two Hudsonian Chickadees (*Parus hudsonicus*) and one Black-poll Warbler (*Dendroica striata*) on the summit of Stratton Mountain in southern Vermont. According to the U. S. Geological Survey the altitude of Stratton Mountain is 3859 feet, and these birds were observed at an altitude of about 3800 feet. The latitude is about 43° 6'—26 miles north of the Massachusetts line. As far as I can learn, the Hudsonian Chickadee has never been recorded from as far south in the breeding season. The Black-poll Warbler has been found by Mr. Bicknell and others in the Catskills, but I can find no other more southern record for it.—FRANCIS H. ALLEN, *West Roxbury, Mass.*